



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

APPLICATION OF THE PELL EQUATION TO THE PROBLEM OF EXTRACTION OF THE CUBE ROOT OF A BINOMIAL SURD.

By E. E. WHITFORD.

In the solution of cubic equations and in certain other cases it is desirable to have a somewhat general method for extracting the cube root of a binomial surd. This may be accomplished by the aid of the Pell Equation. To illustrate by an example let it be required to find the cube root of

$$553090 \sqrt{2} + 53848 \sqrt{211}.$$

Let

$$(553090 \sqrt{2} + 53848 \sqrt{211})^{\frac{1}{3}} = m \sqrt{2} + n \sqrt{211}.$$

Then

$$(553090 \sqrt{2} - 53848 \sqrt{211})^{\frac{1}{3}} = m \sqrt{2} - n \sqrt{211}.$$

Multiplying these two equations,

$$2m^2 - 211n^2 = (611817096200 - 611817098944)^{\frac{1}{3}},$$

$$2m^2 - 211n^2 = (-2744)^{\frac{1}{3}},$$

$$2m^2 - 211n^2 = -14.$$

Let

$$n = 2y.$$

Then

$$m^2 - 422y^2 = -7$$

The solution of this generalized Pell equation may be obtained from one of the convergents of $\sqrt{422}$ when developed into a continued fraction,

$$\sqrt{422} = 20 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{5} \dots}}}$$

or written in another form*

* For tables of continued fractions and for various methods of solving Pell equations see "The Pell Equation," E. E. Whitford, College of the City of New York, 1912.

$$\begin{array}{l} \sqrt{422} = 20, 1, 1, 5, 2, 1, 3, (20) \dots \\ 1, 22, 19, 7, 14, 23, 11, (2) \dots \end{array}$$

where the numbers in the second line denote the values of the form $m^2 - 422y^2$; and hence tell which convergent to select, in this case the fourth. The values of the form are alternately + and —. The convergents are found to be

$$\frac{1}{0}, \frac{20}{1}, \frac{21}{1}, \frac{41}{1}.$$

We select the fourth since for this the value of the form is —7,

$$\begin{array}{l} m = 41. \quad y = 2, \\ n = 4. \end{array}$$

then

$$\therefore (553090\sqrt{2} + 53848\sqrt{211})^{\frac{1}{2}} = 41\sqrt{2} + 4\sqrt{211}.$$

In solving $m^2 - Cy^2 = H$, if $H > 2\sqrt{C}$ certain modifications of the method have to be introduced, but where the solution of the cube root of a binomial surd exists in the form of $m\sqrt{p} \pm n\sqrt{q}$, m, n, p, q , positive integers, this method is always theoretically possible.

Second Illustration:

Extract the cube root of

$$25762\sqrt{2} + 14260\sqrt{7}.$$

(1) Let

$$(25762\sqrt{2} + 14260\sqrt{7}) = m\sqrt{2} + n\sqrt{7};$$

(2) then

$$(25762\sqrt{2} - 14260\sqrt{7})^{\frac{1}{2}} = m\sqrt{2} - n\sqrt{7}.$$

Multiplying (1) and (2)

$$\begin{array}{l} (1327361288 - 1423433200)^{\frac{1}{2}} = 2m^2 - 7n^2; \\ \therefore 2m^2 - 7n^2 = (-96071912)^{\frac{1}{2}}, \\ 2m^2 - 7n^2 = -458. \end{array}$$

Let $n = 2q$.

$$\begin{array}{l} 2m^2 - 28q^2 = -458, \\ (3) \quad m^2 - 14q^2 = -229. \end{array}$$

Now in seeking to solve the equation

$$(A) \quad m^2 - Cq^2 = H, \text{ if } H > 2\sqrt{C},$$

the roots, m, q , can be found from the roots of a similar equation

$$z^2 - Cy^2 = H_1,$$

by the formulas

$$(B) \quad m = \frac{K_1 z + Cy}{H_1}, \quad q = \frac{K_1 y + z}{H_1},$$

provided $K_1 < \frac{1}{2}H$, and such that

$$\frac{K_1^2 - C}{H} = H_1, \text{ an integer.}$$

For substituting the values in (B) into equation (A)

$$\frac{K_1^2 z^2 + 2K_1 C z y + C^2 y^2}{H_1^2} - \frac{CK_1^2 y^2 + 2CK_1 y z + Cz^2}{H_1^2} = H,$$

$$\left(\frac{K_1^2 - C}{H} \right) z^2 - C \left(\frac{K_1^2 - C}{H} \right) y^2 = H_1^2,$$

$$z^2 - Cy^2 = H_1.$$

The roots, z, y , can in their turn be found from the roots of a similar equation

$$l^2 - Ck^2 = H_2,$$

and the process can be repeated until the H_i is less than $2\sqrt{C}$.

Pursuing this method in solving equation (3) we seek

$$K_1 < \frac{1}{2} \cdot 229$$

and such that

$$\frac{K_1^2 - 14}{-229} = H_1$$

and find

$$\frac{2304 - 14}{-229} = -10;$$

$$\therefore K_1 = 48, \quad H_1 = -10.$$

Now applying this method to

$$z^2 - 14y^2 = -10,$$

seeking $K_2 < \frac{1}{2} \cdot 10$, and such that

$$\frac{K_2^2 - 14}{-10} = H_2,$$

we find

$$\frac{4 - 14}{-10} = 1;$$

$$\therefore K_2 = 2, \quad H_2 = 1.$$

But a solution of

$$l^2 - 14k^2 = 1,$$

is evidently $l = 1, \quad k = 0$.

Then by the formulas

$$z = \frac{K_2 l + Ck}{H_2}, \quad y = \frac{K_2 k + l}{H_2};$$

$$z = \frac{2 \cdot 1 + 14 \cdot 0}{1} = 2, \quad y = \frac{2 \cdot 0 + 1}{1} = 1,$$

and by the formulas (B)

$$m = \frac{48 \cdot 2 + 14 \cdot 1}{-10}, \quad q = \frac{48 \cdot 1 + 2}{-10},$$

$$m = 11, \quad q = 5,$$

disregarding quality sign, and since $n = 2q, n = 10$,

$$\therefore (25762\sqrt{2} + 14260\sqrt{7})^{\frac{1}{2}} = 11\sqrt{2} + 10\sqrt{7}.$$

COLLEGE OF CITY OF NEW YORK,
NEW YORK CITY.